

Moving Cellular Materials

as you read

What You'll Learn

- **Describe** the function of a selectively permeable membrane.
- **Explain** how the processes of diffusion and osmosis move molecules in living cells.
- **Explain** how passive transport and active transport differ.

Why It's Important

Cell membranes control the substances that enter and leave the cells in your body.

Review Vocabulary

cytoplasm: constantly moving gel-like mixture inside the cell membrane that contains hereditary material and is the location of most of a cell's life process

New Vocabulary

- passive transport
- diffusion
- equilibrium
- osmosis
- active transport
- endocytosis
- exocytosis

Figure 6 A cell membrane, like a screen, will let some things through more easily than others. Air gets through a screen, but insects are kept out.

Passive Transport

“Close that window. Do you want to let in all the bugs and leaves?” How do you prevent unwanted things from coming through the window? As seen in **Figure 6**, a window screen provides the protection needed to keep unwanted things outside. It also allows some things to pass into or out of the room like air, unpleasant odors, or smoke.


Cells take in food, oxygen, and other substances from their environments. They also release waste materials into their environments. A cell has a membrane around it that works for a cell like a window screen does for a room. A cell's membrane is selectively permeable (PUR mee uh bul). It allows some things to enter or leave the cell while keeping other things outside or inside the cell. The window screen also is selectively permeable based on the size of its openings.

Things can move through a cell membrane in several ways. Which way things move depends on the size of the molecules or particles, the path taken through the membrane, and whether or not energy is used. The movement of substances through the cell membrane without the input of energy is called **passive transport**. Three types of passive transport can occur. The type depends on what is moving through the cell membrane.



Figure 7 Like all other cells in your body, cells in your toes need oxygen.

Describe What is diffusion?



Diffusion Molecules in solids, liquids, and gases move constantly and randomly. You might smell perfume when you sit near or as you walk past someone who is wearing it. This is because perfume molecules randomly move throughout the air. This random movement of molecules from an area where there is relatively more of them into an area where there is relatively fewer of them is called **diffusion**. Diffusion is one type of cellular passive transport. Molecules of a substance will continue to move from one area into another until the relative number of these molecules is equal in the two areas. When this occurs, **equilibrium** is reached and diffusion stops. After equilibrium occurs, it is maintained because molecules continue to move.

Reading Check What is equilibrium?

Every cell in your body uses oxygen. When you breathe, how does oxygen get from your lungs to cells in your big toe? Oxygen is carried throughout your body in your blood by the red blood cells. When your blood is pumped from your heart to your lungs, your red blood cells do not contain much oxygen. However, your lungs have more oxygen molecules than your red blood cells do, so the oxygen molecules diffuse into your red blood cells from your lungs, as shown in **Figure 7**. When the blood reaches your big toe, there are more oxygen molecules in your red blood cells than in your big toe cells. The oxygen diffuses from your red blood cells and into your big toe cells, as shown also in **Figure 7**.

Mini LAB

Observing Molecule Movement

Procedure

1. Use two clean glasses of equal size. Label one *Hot*, then fill it until half full with **very warm water**. Label the other *Cold*, then fill it until half full with **cold water**. **WARNING: Do not use boiling hot water.**
2. Add one drop of **food coloring** to each glass. Carefully release the drop just at the water's surface to avoid splashing the water.
3. Immediately observe the water in the glasses. Record your observations and again after 15 min.

Analysis

What is the relationship between temperature and molecule movement?



Osmosis—The Diffusion of Water Remember that water makes up a large part of living matter. Cells contain water and are surrounded by water. Water molecules move by diffusion into and out of cells. The diffusion of water through a cell membrane is called **osmosis**.

If cells weren't surrounded by water that contains few dissolved substances, water inside of cells would diffuse out of them. This is why water left the carrot cells in this chapter's Launch Lab. Because there were relatively fewer water molecules in the salt solution around the carrot cells than in the carrot cells, water moved out of the cells and into the salt solution.


Losing water from a plant cell causes its cell membrane to come away from its cell wall, as shown on the left in **Figure 8**. This reduces pressure against its cell wall, and a plant cell becomes limp. If the carrot sticks were taken out of salt water and put in pure water, the water around the cells would move into them and they would fill with water. Their cell membranes would press against their cell walls, as shown on the right in **Figure 8**, pressure would increase, and the cells would become firm. That is why the carrot sticks would be crisp again.

Reading Check Why do carrots in salt water become limp?


Osmosis also takes place in animal cells. If animal cells were placed in pure water, they too would swell up. However, animal cells are different from plant cells. Just like an overfilled water balloon, animal cells will burst if too much water enters the cell.

Figure 8 Cells respond to differences between the amount of water inside and outside the cell.

Define What is osmosis?




The carrot stick becomes limp when more water leaves each of its cells than enters them.



Equilibrium occurs when water leaves and enters the cells at the same rate.

Facilitated Diffusion Cells take in many substances. Some substances pass easily through the cell membrane by diffusion. Other substances, such as sugar molecules, are so large that they can enter the cell only with the help of molecules in the cell membrane called transport proteins. This process, a type of passive transport, is known as facilitated diffusion. Have you ever used the drive through at a fast-food restaurant to get your meal? The transport proteins in the cell membrane are like the drive-through window at the restaurant. The window lets you get food out of the restaurant and put money into the restaurant. Similarly, transport proteins are used to move substances into and out of the cell.



Active Transport

Imagine that a football game is over and you leave the stadium. As soon as you get outside of the stadium, you remember that you left your jacket on your seat. Now you have to move against the crowd coming out of the stadium to get back in to get your jacket. Which required more energy—leaving the stadium with the crowd or going back to get your jacket? Something similar to this happens in cells.


Sometimes, a substance is needed inside a cell even though the amount of that substance inside the cell is already greater than the amount outside the cell. For example, root cells require minerals from the soil. The roots of the plant in **Figure 9** already might contain more of those mineral molecules than the surrounding soil does. The tendency is for mineral molecules to move out of the root by diffusion or facilitated diffusion. But they need to move back across the cell membrane and into the cell just like you had to move back into the stadium. When an input of energy is required to move materials through a cell membrane, **active transport** takes place.

Active transport involves transport proteins, just as facilitated diffusion does. In active transport, a transport protein binds with the needed particle and cellular energy is used to move it through the cell membrane. When the particle is released, the transport protein can move another needed particle through the membrane.

Figure 9 Some root cells have extensions called root hairs that may be 5 mm to 8 mm long. Minerals are taken in by active transport through the cell membranes of root hairs.



Transport Proteins Your health depends on transport proteins. Sometimes transport proteins are missing or do not function correctly. What would happen if proteins that transport cholesterol across membranes were missing? Cholesterol is an important lipid used by your cells. Write your ideas in your Science Journal.



Color-enhanced TEM Magnification: 1,400×

Figure 10 One-celled organisms like this egg-shaped one can take in other one-celled organisms using endocytosis.

Endocytosis and Exocytosis

Some molecules and particles are too large to move by diffusion or to use the cell membrane's transport proteins. Large protein molecules and bacteria, for example, can enter a cell when they are surrounded by the cell membrane. The cell membrane folds in on itself, enclosing the item in a sphere called a vesicle. Vesicles are transport and storage structures in a cell's cytoplasm. The sphere pinches off, and the resulting vesicle enters the cytoplasm. A similar thing happens when you poke your finger into a partially inflated balloon. Your finger is surrounded by the balloon in much the same way that the protein molecule is surrounded by the cell membrane. This process of taking substances into a cell by surrounding it with the cell membrane is called **endocytosis** (en duh si TOH sus). Some one-celled organisms, as shown in **Figure 10**, take in food this way.

The contents of a vesicle can be released by a cell using the process called **exocytosis** (ek soh si TOH sus). Exocytosis occurs in the opposite way that endocytosis does. A vesicle's membrane fuses with a cell's membrane, and the vesicle's contents are released. Cells in your stomach use this process to release chemicals that help digest food. The ways that materials can enter or leave a cell are summarized in **Figure 11**.

section 2 review

Summary

Passive Transport

- Cells take in substances and release waste through their cell membranes.
- Facilitated diffusion and osmosis are types of passive transport.

Active Transport

- Transport proteins are involved in active transport.
- Transport proteins can be reused many times.

Endocytosis and Exocytosis

- Vesicles are formed when a cell takes in a substance by endocytosis.
- Contents of a vesicle are released to the outside of a cell by exocytosis.

Self Check

1. **Describe** how cell membranes are selectively permeable.
2. **Compare and contrast** the processes of osmosis and diffusion.
3. **Infer** why endocytosis and exocytosis are important processes to cells.
4. **Think Critically** Why are fresh fruits and vegetables sprinkled with water at produce markets?

Applying Skills

5. **Communicate** Seawater is saltier than tap water. Explain why drinking large amounts of seawater would be dangerous for humans.